

Amoxicillin Fact Sheet

Patient Information: Amoxicillin 500 mg Oral Tablet Amoxicillin Oral Suspension

Take this medicine as prescribed.

Amoxicillin belongs to a class of drugs called penicillin antibiotics. It has been approved by the Food and Drug Administration (FDA) to treat people with infections caused by certain types of bacteria. Amoxicillin has not been approved by the FDA to use when treating people who have been exposed to anthrax. However, if test results show that the anthrax bacteria can be killed by penicillin antibiotics, the use of amoxicillin is recommended to prevent the development of anthrax disease in people who have been exposed to anthrax, when other antibiotics are not as safe to use as with children and pregnant women.

How to take amoxicillin

ADULTS: Take one pill three times a day.

CHILDREN: A child's dose depends on body weight. Give the medicine to your child as directed by the doctor.

Take amoxicillin with a large glass of water. The medication can be taken with or with out food. Taking with food may decrease the chance that upset stomach will occur.

If you miss a dose, start again taking one pill three times a day. Do not take 2 pills to make up for the missed dose. Finish all your pills, even if you feel okay, unless your doctor tells you to stop. If you stop taking this medicine too soon, you may become ill.

For further information, contact the Oklahoma City-County Health Department (405) 425-4437





Amoxicillin Fact Sheet

Side effects

Common side effects of amoxicillin include an upset stomach, vomiting, diarrhea. If you have problems with any of these symptoms, tell your doctor.

Allergic reactions are rare. Signs of an allergic reaction include rash, itching, swelling of the tongue, hands or feet, fever or trouble breathing. If any of these symptoms occur, call your doctor right away.

Precautions

- Be sure to tell the doctor if you are allergic to any medicine
- It is very important to tell your doctor about ALL of the medicine you are currently taking even pills that were bought at the store such as vitamins and antacids.
- Tell your doctor if you have asthma, which is a breathing problem, or any other illness.
- Birth control pills may not work as well when taking this medication. Be sure to use condoms or another form of birth control until you are finished with the entire course of treatment.
- Amoxicillin is safe to take when you are pregnant but be sure your doctor knows if you are pregnant.
- In women, amoxicillin can cause vaginal itching and discharge commonly known as a yeast infection. Tell your doctor if this happens.

